


| Blueberries (Highbush & Lowbush) |

Broad Spectrum Efficacy Against Blueberry Diseases


Quash[®] Fungicide is a highly effective rotational tool that delivers proven, broad spectrum performance for a planned rotation program.

- ▶ **Broad spectrum performance**—highly effective on Mummy berry, fruit rots and more
- ▶ **Preventive protection for when diseases strike**
- ▶ **Helps deliver on optimal yields and crop quality**



Quash delivers excellent, long-lasting mummy berry control.

Multiple applications beginning at green tip through 80–90% bloom; 26 days after last application. Rates were for research purposes. Source: North Carolina State University



Quash delivers outstanding mummy berry control.

Source: Oregon State University

Quash Efficacy vs. Other FRAC Group 3 Mode-of-Action Fungicides

	<i>Quash</i> (metconazole)	Indar* (fenbuconazole)	Proline [®] (prothioconazole)
	PHI = 7 Days	PHI = 30 Days	PHI = 30 Days
Mummy Berry	E	E	E
Ripe Rot (Anthracnose)	E	NA*	UN
Anthracnose Leaf Spot	E	E	UN
Phomopsis Twig Blight	E	E	E
Septoria Leaf Spot	E	E	G
Rust	VG	G	E
Alternaria Rot	E	NA	NA

*Tank mix with Captan products during bloom to prevent rots.

2023 Southeast Regional Blueberry Integrated Management Guide efficacy ratings: E = excellent, VG = very good, G = good, F = fair, P = poor, NA = not recommended, UN = control unknown. These ratings are benchmarks, actual performance will vary.

How To Use

Rate	2.5 oz/A per application (maximum 7.5 oz/A per year) <ul style="list-style-type: none"> Up to 3 foliar applications per season Do not make more than 2 sequential applications
Timing	When conditions favor disease development and prior to infection; apply on a 7–14 day interval
Method	Ground; air
Spray Volume	20 GPA by ground; 10 GPA by air; make sure volume is sufficient to obtain coverage of blossoms, foliage and/or fruit
REI / PHI	12 hours / 7 days

Other Important Information

- ▶ Use *Quash* as part of an Integrated Pest Management (IPM) program
- ▶ Rotate with a non-Group 3 fungicide that is active on the target disease

Maximum Residue Limits (MRLs)

Countries with Harmonized Highbush and Lowbush Blueberry MRLs for Metconazole (the Active Ingredient in <i>Quash</i>)			
Canada	Japan	South Korea	Taiwan
0.4	0.4	2.0	0.4

All MRLs in table are in parts per million (ppm).



Products That Work, From People Who Care® | valent.com | 800-6-VALENT (682-5368)

Always read and follow label instructions.

Quash and *Products That Work, From People Who Care* are registered trademarks of Valent U.S.A. LLC. Indar is a registered trademark of Dow AgroSciences. LLC. Pristine is a registered trademark of BASF. Proline is a registered trademark of Bayer Group. Tilt is a registered trademark of a Syngenta Group Company. ©2024 Valent U.S.A. LLC. All rights reserved. Printed in the U.S.A.
2024-QUA-8014 7/24



LEARN MORE